

Take 5 for Safety


E. Lessard
June 17, 2014

BROOKHAVEN
NATIONAL LABORATORY

a passion for discovery




How to minimize Lyme disease risk




- Outdoors at BNL is a high-risk tick area and you should take precautions:
- Wear enclosed shoes or boots, long-sleeved shirts, and long pants
- If working outdoors, tuck your pant legs into your shoes or boots to prevent ticks from crawling up your legs
- Use an insect repellent containing 20% to 30% DEET (N,N-diethyl-met-toluamide)
- Wear light-colored clothing to help you see ticks more easily
- Keep long hair pulled back or wear a cap for protection
- Don't sit on the ground outside
- Check yourself for ticks regularly – both indoors and outdoors

Tritium at BLIP

- Three wells downstream of BLIP are all below Drinking Water Standard (DWS) for tritium; <20,000 pCi/L
- Caps are working




Tritium at AGS



- Wells downstream of E20, J10, U target area and U dump are all below DWS for tritium; <20,000 pCi/L
- Wells downstream of V are slightly above or below the DWS; down from 4E6 pCi/L
- Caps are working


Tritium at RHIC




- Wells downstream of beam dumps are all below DWS for tritium; <20,000 pCi/L
- Wells downstream of collimators are all below DWS for tritium; <20,000 pCi/L

BNL Injury Rates and Injuries

Injury Rates by Fiscal Year



FY2014 Injuries by Directorate



- A number of injuries occur during work planning and clean-up
- Distracted walking may be a precursor to some injuries

Distracted Walking

Ohio Data

